

A Nonperturbative Construction of Retarded Off-Shell Intertwiners of Covariant Phase Spaces in Classical Field Theory

Pedro Lauridsen Ribeiro
pedro.lauridsen.ribeiro@desy.de

II. Institut für theoretische Physik – Universität Hamburg¹

(Joint work with R. Brunetti (Trento – brunetti@science.unitn.it) and K. Fredenhagen (Hamburg – klaus.fredenhagen@desy.de))

25.III.2009

¹This work is being performed under the aegis of the Collaborative Research Centre 676 – “Particles, Strings and the Early Universe – The Structure of Matter and Spacetime”.

- 1 The stage
 - Motivation and Setup
 - Further properties
- 2 The actors
- 3 Strategy towards a proof
 - Nash–Moser–Hörmander theorem
 - Tame differentiability
 - End of proof
- 4 Consequences
- 5 Conclusions and perspectives

Motivation and Setup

In this talk, we shall be interested in the following problem: consider (for concreteness; watch out for Final Considerations)...

- A scalar field $\phi \in \mathcal{C}^\infty(\mathcal{M})$ in a globally hyperbolic spacetime (\mathcal{M}, g) , and
- Two (1st-order) action functionals

$$S_i[\phi] = \int_{\mathcal{M}} \sqrt{|\det g(x)|} dx \mathcal{L}_i(x, \phi(x), \partial^1 \phi(x)), \quad i = 1, 2$$

with (semilinear, strictly hyperbolic) Euler-Lagrange derivatives

$$S_{i(1)}[\phi] = \nabla_a \frac{\partial \mathcal{L}_i}{\partial \nabla_a \phi} - \frac{\partial \mathcal{L}_i}{\partial \phi},$$

such that

- i.) $S_2 \doteq S$ is quadratic (“free”),
 ii.)

$$S_1 - S_2 = \lambda F(h) = \lambda \int_{\mathcal{M}} \sqrt{|\det g(x)|} dx h(x) \mathcal{L}_{int}(x, \phi(x), \partial^1 \phi(x))$$

with $h \in \mathcal{C}_c^\infty(\mathcal{M})$ (“spacetime-cutoff” interaction term), $\lambda > 0$, and

- iii.) $F(h)_{(1)}[\phi]$ depends pointwise on ϕ and at most its first derivatives $\nabla \phi$.

We want to

Main Goal & Definition

Prove the existence of a map $r_{S_1, S_2} : \mathcal{C}^\infty(\mathcal{M}) \rightarrow \mathcal{C}^\infty(\mathcal{M})$ such that

$$S_{1(1)} \circ r_{S_1, S_2} = S_{2(1)}, \tag{1}$$

$$r_{S_1, S_2}(\phi)(x) = \phi(x), x \notin J^+(\text{supp} h). \tag{2}$$

We call r_{S_1, S_2} the **retarded Møller operator** of S_1 w.r.t. S_2 .

- r_{S_1, S_2} appears naturally in the context of perturbative algebraic QFT (Dütsch–Fredenhagen CMP'03, Brunetti–Fredenhagen arXiv:0901.2063, Brunetti–Dütsch–Fredenhagen arXiv:0901.2038), where \hbar plays **both** the role of an **IR regulator** and of a **localization** for the algebra of perturbative interacting fields.
- When acting on solutions of $S_{2(1)}[\phi] = 0$ r_{S_1, S_2} can be seen as an **intertwiner** of (on-shell) covariant phase spaces or, equivalently, as the solution of a **“covariant” Cauchy problem**.
- (1)–(2) also mean that $r_{S_1, S_2}(\phi)$ solves an **inhomogeneous** (off-shell) nonlinear hyperbolic PDE with prescribed initial conditions in the past of $\text{supp } \hbar \Rightarrow$ **very few** rigorous well-posedness results exist, qualitative behaviour of solutions can be **dramatically changed** – (parabolic) example: (incompressible) **Navier-Stokes equations**

$$\underbrace{\partial_t \mathbf{v} - \nu \Delta \mathbf{v}}_{\text{free (heat) part}} + \underbrace{(\mathbf{v} \cdot \nabla) \mathbf{v}}_{\text{interaction}} = \underbrace{-\nabla p}_{\text{source}} \rightarrow \begin{cases} \nabla p = 0 : \text{only laminar flow} \\ \nabla p \neq 0 : \text{turbulence!} \end{cases}$$

Coupling as an off-shell flow parameter \Rightarrow Main Claim

- It's clear that r_{S_1, S_2} exist on shell whenever local well posedness for $S_{(1)}[\psi] = (S + \lambda F(h))_{(1)}[\psi] = 0$ in a nbg. of $\text{supp} h$ holds. More in general, in the future of $\text{supp} h$ (1) tells us that $\psi = r_{S_1, S_2}(\phi) - \phi$ solves $S_{(1)}[\psi] = 0 \Rightarrow$ finding r_{S_1, S_2} boils down to finding it **locally!**
- Before we start...

Caveat 1

Notice that we cannot use (2) and apply the (background-independent) **retarded fundamental solution** Δ_S^R of $S^{(1)} = S^{(2)}[\psi], \forall \psi$ to (1) directly to obtain the so-called **Yang-Feldman equation**

$$r_{S+\lambda F(h), S}(\phi) = \phi - \lambda \Delta_S^R \circ F(h)_{(1)}[r_{S+\lambda F(h), S}(\phi)],$$

for $S_{(1)}[\psi]$ doesn't necessarily have compact support or even enough decay at infinity!

- Differentiating (1) w.r.t. λ leads to

$$(S_{(1)} + \lambda F(h)_{(1)})^{(1)} [r_{S+\lambda F(h), S}(\phi)] \circ \frac{d}{d\lambda} r_{S+\lambda F(h), S}(\phi) + F_{(1)} [r_{S+\lambda F(h), S}(\phi)] = 0. \quad (3)$$

- Now we invoke (2) and apply the retarded fundamental solution $\Delta_{S+\lambda F(h)}^R [r_{S+\lambda F(h), S}(\phi)]$ of the **linearised Euler-Lagrange operator** $(S + \lambda F(h))_{(1)}^{(1)} [r_{S+\lambda F(h), S}(\phi)]$ around the background $r_{S+\lambda F(h), S}(\phi)$ to the left of both sides of (3):

$$\frac{d}{d\lambda} r_{S+\lambda F(h), S}(\phi) = -\Delta_{S+\lambda F(h)}^R [r_{S+\lambda F(h), S}(\phi)] \circ F_{(1)}(h) [r_{S+\lambda F(h), S}(\phi)], \quad (4)$$

which shows that $\psi(\lambda) \doteq r_{S+\lambda F(h), S}(\phi)$ is the unique solution of the **flow equation** (4) with initial condition $\psi(0) = \phi$.

- Formally integrating (4) w.r.t. λ on both sides and using the initial condition above, we arrive at

$$r_{S+\lambda F(h),S}(\phi) = \phi - \int_0^\lambda d\lambda' \Delta_{S+\lambda' F(h)}^R [r_{S+\lambda' F(h),S}(\phi)] \circ F_{(1)}(h) [r_{S+\lambda' F(h),S}(\phi)]. \quad (5)$$

- We could keep proceeding formally by iterating (4) and write $r_{S+\lambda F(h),S}(\phi)$ as a formal power series [Dütsch–Fredenhagen *ibid.*]

$$r_{S+\lambda F(h),S} \sim \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} \frac{d^k}{d\lambda^k} \Big|_{\lambda=0} r_{S+\lambda F(h),S}.$$

This is actually the method used in (perturbative algebraic) quantum field theory when performed after composition with functionals, barring renormalisation issues.

- However, our aim is **nonperturbative**, and thus achieved by looking at the map

$$\psi(\lambda) \mapsto \phi(\lambda) = \psi(\lambda) + \int_0^\lambda d\lambda' \Delta_{S+\lambda'F(h)}^R[\psi(\lambda')] \circ F_{(1)}(h)[\psi(\lambda')], \quad (6)$$

which just defines the **inverse** $r_{S+\lambda F(h),S}^{-1}$ of $r_{S+\lambda F(h),S}$.

- From now on, for the sake of pedagogy we shall set $(\mathcal{M}, \mathbf{g}) = \mathbb{R}^{1,d-1} \ni (x^0 = t, \mathbf{x})$.

Main Claim

The map (6) is invertible in a neighbourhood of zero in $\mathcal{C}^1([0, \Lambda], \mathcal{C}^\infty(\mathcal{M}))$; its inverse satisfies (1), (2).

Towards a proof of Main Claim

- The crucial step in our proof is to obtain a priori estimates on $\Delta_{S+\lambda F(h)}^R[\psi]$ in terms of **both** the linear and the nonlinear (background) arguments. These are essentially **refined energy estimates for $S_{(1)}^{(1)} + \lambda F(h)_{(1)}^{(1)}$** which state explicitly their dependence on the latter's coefficients, and were originally obtained by Klainerman [Klainerman '78–'80–'82].
- Suppose that there exist $0 < T$ such that $\text{supp} h$ is contained in the interior of the slab $\{(t, x) : 0 \leq t \leq T\}$, and define the **energy norms**

$$\|\psi\|_{E^k} \doteq \sup_{t' \in [0, T]} \|\psi(t', \cdot)\|_{H_x^{(k+1)}} + \sup_{t' \in [0, T]} \|\partial_t \psi(t', \cdot)\|_{H_x^{(k)}}.$$

Caveat 2

It feels tempting to apply a fixed-point strategy to (5), but the Proposition below shows that $r_{S+\lambda F(h), S}(\phi) - \phi$ is **not** a Lipschitz map with respect to any $\|\cdot\|_{E^k}$! (PDE folk wisdom: **"hyperbolic equations are not strongly stable w.r.t. perturbations of the coefficients."** [Tataru ICM '02])

Proposition (refined energy estimates)

For $\phi, \delta\phi \in E^\infty \doteq \{\psi : \|\psi\|_{E^k} < +\infty, \forall k \geq 0\}$ we have

$$\|\Delta_{S+\lambda F(h)}^R[\phi]\delta\phi\|_{E^0} \leq D_0 \sup_{t' \in [0, T]} \|\delta\phi\|_{L_x^2}, \quad (7)$$

$$\|\Delta_{S+\lambda F(h)}^R[\phi]\delta\phi\|_{E^k} \leq D_k \left(\|\delta\phi\|_{E^{k-1}} + \sup_{t' \in [0, T]} |(hF_{(1)}^{(1)})(t', \cdot)|_{\mathcal{C}_x^k} \|\delta\phi\|_{E^0} \right), \quad k \geq 1, \quad (8)$$

where $D_k, k \geq 0$ are constants which depend **only on d, T and $\|\phi\|_{\mathcal{C}^1(\text{supph})}$** .

- Applying Sobolev inequalities and Schauder estimates to the spatial \mathcal{C}^k norms of $(hF_{(1)}^{(1)})(t', \cdot)$ in (8), we arrive at

$$\|\Delta_{S+\lambda F(h)}^R[\phi]\delta\phi\|_{E^k} \leq D'_k [\|\delta\phi\|_{E^{k-1}} + (1 + \|\phi\|_{E^{k+1+[\frac{d+1}{2}]}}) \|\delta\phi\|_{E^0}], \quad (9)$$

where $[s]$ gives the integer part of s .

Nash–Moser–Hörmander iteration scheme

- The argument above shows that **one loses $1 + \lfloor \frac{d+1}{2} \rfloor$ derivatives** at each iteration when trying to solve (5) by a fixed-point method (see Caveat 2!). This phenomenon has **no on-shell counterpart**.
- Alternative: use a **Newton iteration scheme** \Rightarrow if it converges, it does so superexponentially; not the case here, again due to loss of derivatives. This can be fixed by applying suitable smoothing operators that make a “multiscale” decomposition of momentum space at each iteration step. The result is the celebrated

Theorem (Nash–Moser–Hörmander)

Let $\Phi : \mathcal{U} \subseteq E^\infty \cap \{\psi : \|\psi - \psi_0\|_{E^\mu} < R\} \rightarrow E^\infty$, $\mu \in \bar{\mathbb{Z}}_+$, $R > 0$ be twice Gâteaux differentiable satisfying for all $k \geq 0$ the **tame estimates**

$$\|\Phi(\psi)\|_{E^k} \leq C_k(1 + \|\psi\|_{E^{k+r_0}}) \text{ for some } r_0 > 0, \quad (10)$$

$$\|\Phi'(\psi)(\delta\psi)\|_{E^k} \leq C'_k[(1 + \|\psi\|_{E^{k+r_1}})\|\delta\psi\|_{E^{s_1}} + \|\delta\psi\|_{E^{k+s_1}}] \text{ for some } r_1, s_1 > 0, \quad (11)$$

$$\begin{aligned} \|\Phi''(\psi)(\delta_1\psi, \delta_2\psi)\|_{E^k} \leq C_k'' [(1 + \|\psi\|_{E^{k+r_2}}) \|\delta_1\psi\|_{E^{s_2}} \|\delta_2\psi\|_{E^{t_2}} + \|\delta_1\psi\|_{E^{s_2}} \|\delta_2\psi\|_{E^{k+t_2}} \\ + \|\delta_1\psi\|_{E^{k+t_2}} \|\delta_2\psi\|_{E^{s_2}}], \text{ for some } r_2, s_2, t_2 > 0, \end{aligned} \quad (12)$$

and such that for all ψ in $\mathcal{V} \subset \{\psi : \|\psi - \psi_0\|_{E^{\mu'}} < R'\}$, $\mu' \in \bar{\mathbb{Z}}_+$, $R' > 0$ there is a right inverse $\Psi(\psi)$ to $\Phi'(\psi)$ w.r.t. the linear factor satisfying for all $k \geq 0$ the tame estimates

$$\|\Psi'(\psi)(\delta\psi)\|_{E^k} \leq C_k''' [(1 + \|\psi\|_{E^{k+a_1}}) \|\delta\psi\|_{E^{b_1}} + \|\delta\psi\|_{E^{k+b_1}}] \text{ for some } a_1, b_1 > 0. \quad (13)$$

Then, for all k sufficiently large, there is a $R_k > 0$ such that for all $\phi \in E^\infty$ fulfilling $\|\phi\|_{E^{k+b_1}} < R_k$ the equation $\Phi(\psi) = \Phi(\psi_0) + \phi$ has a unique solution $\psi = \psi(\phi)$ such that $\|\psi(\phi) - \psi_0\|_{E^k} \leq R'' \|\phi\|_{E^{k+b_1}}$. In particular, if ϕ also belongs to E^∞ , so does $\psi(\phi)$.

In our problem, we take $\psi_0 \equiv 0$ and add a dependence in λ .

Tame (Gâteaux) differentiability of $\Delta_{S+\lambda F(h)}^R[\psi]$

To check that Φ_λ fulfills the hypotheses of the Theorem, first we collect some following formulae coming directly from the definition of a fundamental solution [Dütsch–Fredenhagen *ibid.*]:

$$\Delta_{S+\lambda F(h)}^{R(1)}[\psi](\delta\psi) = -\Delta_{S+\lambda F(h)}^R[\psi] \circ F(h)_{(1)}^{(2)}[\psi](\delta\psi, \Delta_{S+\lambda F(h)}^R[\psi]), \quad (14)$$

$$\frac{d}{d\lambda} \Delta_{S+\lambda F(h)}^R[\psi] = -\Delta_{S+\lambda F(h)}^R[\psi] \circ F(h)_{(1)}^{(1)}[\psi] \circ \Delta_{S+\lambda F(h)}^R[\psi], \quad (15)$$

$$\begin{aligned} & \Delta_{S+\lambda F(h)}^{R(2)}[\psi](\delta_1\psi, \delta_2\psi) = \\ &= \Delta_{S+\lambda F(h)}^R[\psi] \circ F(h)_{(1)}^{(2)}(\delta_1\psi, \Delta_{S+\lambda F(h)}^R[\psi]) \circ F(h)_{(1)}^{(2)}(\delta_2\psi, \Delta_{S+\lambda F(h)}^R[\psi]) + \\ &+ \Delta_{S+\lambda F(h)}^R[\psi] \circ F(h)_{(1)}^{(2)}(\delta_2\psi, \Delta_{S+\lambda F(h)}^R[\psi]) \circ F(h)_{(1)}^{(2)}(\delta_1\psi, \Delta_{S+\lambda F(h)}^R[\psi]) + \\ &- \Delta_{S+\lambda F(h)}^R[\psi] \circ F(h)_{(1)}^{(3)}(\delta_1\psi, \delta_2\psi, \Delta_{S+\lambda F(h)}^R[\psi]). \end{aligned} \quad (16)$$

Equation (15) shows in particular that $\Delta_{S+\lambda F(h)}^R[\psi]$ is strongly differentiable (hence strongly continuous) in λ , thus allowing all the computations we need.

Tame estimates for iteration map, end of proof

From (14) and (16), one get the following formulae for the first two derivatives of the iteration map Φ_λ (6):

$$\begin{aligned} \Phi'_\lambda(\psi)(\delta\psi) &= \delta\psi + \\ &+ \int_0^\lambda d\lambda' \left(\Delta_{S+\lambda'F(h)}^R[\psi] \circ F(h)_{(1)}^{(1)}[\psi](\delta\psi) + \Delta_{S+\lambda'F(h)}^{R(1)}[\psi](\delta\psi) \circ F(h)_{(1)}[\psi] \right), \end{aligned} \quad (17)$$

$$\begin{aligned} \Phi''_\lambda(\psi)(\delta_1\psi, \delta_2\psi) &= \int_0^\lambda d\lambda' \left(\Delta_{S+\lambda'F(h)}^R[\psi] \circ F(h)_{(1)}^{(2)}[\phi](\delta_1\phi, \delta_2\phi) + \right. \\ &+ \Delta_{S+\lambda'F(h)}^{R(1)}[\psi](\delta_1\psi) \circ F(h)_{(1)}^{(1)}[\phi](\delta_2\phi) + \Delta_{S+\lambda'F(h)}^{R(1)}[\psi](\delta_2\psi) \circ F(h)_{(1)}^{(1)}[\phi](\delta_1\phi) + \\ &\left. + \Delta_{S+\lambda'F(h)}^{R(2)}[\psi](\delta_1\psi, \delta_2\psi) \circ F(h)_{(1)}[\phi] \right), \end{aligned} \quad (18)$$

where $\Delta_{S+\lambda'F(h)}^{R(1)}[\psi]$ and $\Delta_{S+\lambda'F(h)}^{R(2)}[\psi]$ are respectively given by (14) and (16). Notice that $\frac{d}{d\lambda}\Phi'_\lambda(\psi)$, seen as a linear map acting on $\delta\psi$ for fixed ψ , **doesn't lose derivatives**, due to the fact that the assumed loss in $F(h)_{(1)}$ is exactly compensated by the **smoothing effect** of $\Delta_{S+\lambda'F(h)}^R[\psi]$.

- The Proposition, together with Schauder estimates, show that Φ_λ satisfy the tame estimate (10) with $a_0 = \lfloor \frac{d+1}{2} \rfloor + 1$ for $\sup_{\lambda' \in [0, \lambda]} \|\psi(\lambda')\|_{E^{\lfloor \frac{d+1}{2} \rfloor + 1}} < R$, that is, $\mu = \lfloor \frac{d+1}{2} \rfloor + 1$.
- Formulae (17)–(18) show that $\Phi'_\lambda(\psi)(\delta\psi)$ and $\Phi''_\lambda(\psi)(\delta_1\psi, \delta_2\psi)$ fulfill resp. the tame estimates (11) and (12) with $r_1 = r_2 = \lfloor \frac{d+1}{2} \rfloor + 1$ and $s_1 = s_2 = t_2 = 1$.
- Finally, due to (15) and the remark following (18), $\frac{d}{d\lambda}\Phi'_\lambda(\psi)$ is a **bounded** and **uniformly strongly continuous** (in λ) linear map $\Rightarrow \Phi'_\lambda(\psi)$ be inverted by means of a Dyson series. Iterating the tame estimate for $\frac{d}{d\lambda}\Phi'_\lambda(\psi)$, together with the argument for the convergence for the Dyson series, leads to the tame estimate (13) with $a_1 = \lfloor \frac{d+1}{2} \rfloor + 1$, $b_1 = 1$ and $\mu' = \lfloor \frac{d+1}{2} \rfloor + 2$ for the right inverse.
- Now... Just **plug in** the data above, **run** the “Nash–Moser–Hörmander machine”, and we get **local existence and uniqueness of $r_{S+\lambda F(h), S}$ in E^∞** . The intertwining relation (1) shows that actually **$r_{S+\lambda F(h), S}(\phi) \in \mathcal{C}^\infty$ for $\phi \in \mathcal{C}^\infty$** . □

Structural consequences

The existence and properties of $r_{S+\lambda F(h),S}$ have fundamental implications for the underlying **Poisson structure** of any classical field theory determined by an action functional S , given by the **Peierls bracket**

$$\{F, G\}_S \doteq F_{(1)}[\cdot] \circ (\Delta_S^R[\cdot] - \Delta_S^A[\cdot]) \circ G_{(1)}[\cdot]$$

of local functionals F, G . Here $\Delta_S^A[\psi]$ is the **advanced fundamental solution** of $S_{(1)}^{(1)}[\psi]$ around the background ψ , which is simply the **adjoint** of $\Delta_S^R[\psi]$.

Corollary $r_{S+\lambda F(h),S}$ is a **canonical transformation**, i.e. it intertwines the Poisson structures associated to S and $S + \lambda F(h)$:

$$\{\cdot, \cdot\}_{S+\lambda F(h)} \circ r_{S+\lambda F(h)} = \{\cdot, \cdot\}_S.$$

In particular, even off shell does it allow one to put $\{\cdot, \cdot\}_{S+\lambda F(h)}$ in **“normal form”**, i.e. to make it **locally background-independent** (“**Functional Darboux Theorem**”).

Scholium The space of local functionals vanishing on solutions of $(S + \lambda F(h))_{(1)}[\psi] = 0$ is a **Poisson ideal**.

Coda: final considerations

- We've shown the existence of r_{S_1, S_2} for “sufficiently small” field configurations around a given one. This latter condition can be controlled in general by adjusting λ (**coupling strength**) or supph (**lifespan**).
- If the Cauchy problem for $S_{1(1)}[\psi] = 0$ is well-posed **in the large**, one can use (2) and the **composition property** of r_{S_1, S_2}

$$r_{S, S} = \mathbb{1}, r_{S_2, S_3} \circ r_{S_1, S_2} = r_{S_1, S_3}$$

stemming from (1) to **remove the cutoff** (i.e. dependence on h) \Rightarrow probably impossible off shell, unless probably in a suitable algebraic sense (“algebraic adiabatic limit” – [Brunetti–Dütsch–Fredenhagen *ibid.*]).

- We illustrated our strategy for the case of a scalar field in $\mathbb{R}^{1,d-1}$, but the argument carries through for **arbitrary sections** in **any globally hyperbolic spacetime** \Rightarrow one has a **local energy estimate** of the same form as (7)–(8) by combining Klainerman's argument with the estimates in [Hawking–Ellis '73]; only the control of the extra error terms due to **curvature** and the **absence of Killing fields** is more cumbersome.
- Alas, the more general **quasilinear** case (e.g. general relativity) seems to pose some **new difficulties**; the Dyson series argument to invert Φ'_λ w.r.t. the linear factor fails since one then loses one derivative at each order. It seems to be possible, however, to circumvent this issue by means of **paradifferential calculus** (PLR, work in progress).